

STROKE OCCURS IN TODDLERS, CHILDREN, AND TEENS. STROKE IS AN EMERGENCY AT ANY AGE!

WARNING SIGNS



IF ONE OR MORE OF THESE SIGNS ARE PRESENT, DON'T DELAY!

CALL 911 OR THE EMERGENCY SYSTEM
IN YOUR COUNTRY!

ADDITIONAL WARNING SIGNS IN CHILDREN

- Severe sudden headache, especially with vomiting and sleepiness
- Sudden weakness or numbness on one side of the body (face, arm and/or leg)
- Sudden confusion, difficulty speaking or understanding others
- Sudden trouble seeing to one side or loss of vision
- Sudden difficulty walking, dizziness, loss of balance or coordination
- New-onset of seizures usually on one side of the body

IF IT LOOKS LIKE A STROKE

THINK STROKE

Signs of stroke are often missed in children and teens because there is a lack of awareness that strokes can happen in this age group.



About HALF of the children presenting with a stroke have a PREVIOUSLY IDENTIFIED RISK FACTOR

10

Stroke is one of the TOP 10 CAUSES OF DEATH in children



The use of tPA (a clot busting treatment) or clot retrieval devices are

NOT FDA APPROVED
FOR USE IN CHILDREN
in the U.S.

MEDICAL CONDITIONS ASSOCIATED WITH STROKE INCLUDE:

Sickle Cell Disease, Moyamoya Syndrome, Arterial Dissection, Autoimmune Disorders, Congenital Heart Disease, Blood Clotting Disorders

Learn more at iapediatricstroke.org and strokeassociation.org

Sources

iapediatricstroke.org/infographic.pdf iapediatricstroke.org/2014%20AHA.fact.pdf heart.org/idc/groups/heart-public/@wcm/@sop/@smd/documents/downloadable/ucm_472920.pdf

International Alliance